

Epidemiology, Diagnosis, and Treatment of Neck Pain

Steven P. Cohen, MD

From the Department of Anesthesiology and Critical Care Medicine and Department of Physical Medicine and Rehabilitation, Johns Hopkins School of Medicine, Baltimore, MD; and Department of Anesthesiology and Department of Physical Medicine and Rehabilitation, Uniformed Services University of the Health Sciences, Bethesda, MD.

CME Activity

Target Audience: The target audience for *Mayo Clinic Proceedings* is primarily internal medicine physicians and other clinicians who wish to advance their current knowledge of clinical medicine and who wish to stay abreast of advances in medical research.

Statement of Need: General internists and primary care physicians must maintain an extensive knowledge base on a wide variety of topics covering all body systems as well as common and uncommon disorders. *Mayo Clinic Proceedings* aims to leverage the expertise of its authors to help physicians understand best practices in diagnosis and management of conditions encountered in the clinical setting.

Accreditation: Mayo Clinic College of Medicine is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

Credit Statement: Mayo Clinic College of Medicine designates this journal-based CME activity for a maximum of 1.0 *AMA PRA Category 1 Credit(s)*.^{TM1} Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Learning Objectives: On completion of this article, you should (1) be able to distinguish the different types (eg, neuropathic or nociceptive) of neck pain, (2) be able to identify "red flags" that may warrant advanced work-up, (3) be familiar with the risk factors for development of neck pain and its natural course, (4) know when and in whom advanced diagnostic testing may be helpful, and (5) be able to identify which patients to refer for specialty care (eg, injections or surgery).

Disclosures: As a provider accredited by ACCME, Mayo Clinic College of Medicine (Mayo School of Continuous Professional Development) must ensure balance, independence, objectivity, and scientific rigor in its educational activities. Course Director(s), Planning Committee members, Faculty, and all others who are in a position to control the content of this educational activity are required to disclose all relevant financial relationships with any commercial interest related to the subject matter of the educational activity.

Safeguards against commercial bias have been put in place. Faculty also will disclose any off-label and/or investigational use of pharmaceuticals or instruments discussed in their presentation. Disclosure of this information will be published in course materials so that those participants in the activity may formulate their own judgments regarding the presentation.

In their editorial and administrative roles, William L. Lanier, Jr, MD, Terry L. Jopke, Kimberly D. Sankey, and Nicki M. Smith, MPA, have control of the content of this program but have no relevant financial relationship(s) with industry. Dr Cohen serves as a consultant for Regenesys Biomedical and Semnur Pharmaceuticals, Inc, and is on the advisory board of Kimberly Clark Health Care. This work was funded in part by a Congressional grant from the Center for Rehabilitation Sciences Research. The role of the funding source was only to pay for salary support for the author.

Method of Participation: In order to claim credit, participants must complete the following:

1. Read the activity.
2. Complete the online CME Test and Evaluation. Participants must achieve a score of 80% on the CME Test. One retake is allowed. Visit www.mayoclinicproceedings.com, select CME, and then select CME articles to locate this article online to access the online process. On successful completion of the online test and evaluation, you can instantly download and print your certificate of credit.

Estimated Time: The estimated time to complete each article is approximately 1 hour.

Hardware/Software: PC or MAC with Internet access.

Date of Release: 02/01/2015

Expiration Date: 01/31/2017 (Credit can no longer be offered after it has passed the expiration date.)

Privacy Policy: <http://www.mayoclinic.org/global/privacy.html>

Questions? Contact dletsupport@mayo.edu.

Abstract

Neck pain is the fourth leading cause of disability, with an annual prevalence rate exceeding 30%. Most episodes of acute neck pain will resolve with or without treatment, but nearly 50% of individuals will continue to experience some degree of pain or frequent occurrences. History and physical examination can provide important clues as to whether the pain is neuropathic or mechanical and can also be used to identify "red flags" that may signify serious pathology, such as myelopathy, atlantoaxial subluxation, and metastases. Magnetic resonance imaging is characterized by a high prevalence of abnormal findings in asymptomatic individuals but should be considered for cases involving focal neurologic symptoms, pain refractory to conventional treatment, and when referring a patient for interventional treatment. Few clinical trials have evaluated treatments for neck pain. Exercise treatment appears to be beneficial in patients with neck pain. There is some evidence to support muscle relaxants in acute neck pain associated with muscle spasm, conflicting evidence for epidural corticosteroid injections for radiculopathy, and weak positive evidence for cervical facet joint radiofrequency denervation. In patients with radiculopathy or myelopathy, surgery appears to be more effective than nonsurgical therapy in the short term but not in the long term for most people.

© 2015 Mayo Foundation for Medical Education and Research ■ *Mayo Clin Proc.* 2015;90(2):284-299

In the past few years, several reviews have been written on neck pain, although far fewer than on back pain, which often, but not always, involves similar mechanisms. Most of these reviews have targeted a specific

specialty audience and have focused on one particular aspect of neck pain, rather than encompassing a broad overview aimed toward a general medical audience. The purpose of this review is to provide such an article, to

include epidemiological aspects, classification, the natural course of neck pain, and an evidence-based, comprehensive guide to work-up, diagnosis, and treatment.

METHODS

Databases on Medline via PubMed and Ovid, Embase, and the Cochrane Database of Systematic Reviews were searched using the key words *neck pain*, *cervical pain*, *cervical radiculopathy*, and *cervical myelopathy*, with no date restrictions. For individual sections, key words relating to specific topics (eg, *physical exam*, *history*, *radiological*, *surgery*, *epidural steroid injection*, *antidepressant*, *spinal manipulation*, *acupuncture*, *complementary and alternative medicine*) were identified and cross-referenced with the initial search terms using the aforementioned databases. Prime references and additional articles were obtained by cross-referencing all search terms with *review article* and manually searching through reference lists.

OVERVIEW AND EPIDEMIOLOGY

The physical, psychological, and socioeconomic impact of neck pain is underappreciated. According to the Global Burden of Disease 2010 Study, neck pain is the fourth leading cause of years lost to disability, ranking behind back pain, depression, and arthralgias.¹ Approximately half of all individuals will experience a clinically important neck pain episode over the course of their lifetime.² There is substantial heterogeneity in the reported prevalence rates of neck pain; however, most epidemiological studies report an annual prevalence ranging between 15% and 50%,²⁻⁵ with one systematic review reporting a mean rate of 37.2%.² The prevalence of neck pain is higher in females and peaks in middle age.²⁻⁵ Neck pain is associated with several comorbidities including headache, back pain, arthralgias, and depression.^{3,5}

Who Gets Neck Pain?

The factors associated with the development and persistence of neck pain overlap considerably with those of other musculoskeletal conditions. The prevalence of neck pain is higher in females than in males, and the literature is mixed as to whether it peaks or plateaus in middle age.²⁻⁶ Variables associated with neck pain that overlap with other rheumatologic

conditions include genetics, psychopathology (eg, depression, anxiety, poor coping skills, somatization), sleep disorders, smoking, and sedentary lifestyle. For obesity, the results of epidemiological studies have usually but not always found a positive association between neck and shoulder pain and body mass index.^{3,6-9} Some of the reasons why obese individuals may be predisposed to neck pain include elevated systemic inflammation, deleterious structural changes, increased mechanical stress and ground reaction force, diminished muscle strength, more psychosocial issues, and greater disability related to kinesiophobia compared with nonoverweight people.¹⁰ Unique risk factors for neck pain include trauma (eg, traumatic brain and whiplash injuries) and certain sports injuries (eg, wrestling, ice hockey, football). Although certain occupations such as office and computer workers, manual laborers, and health care workers, have been found in some studies to have a higher incidence of neck pain, the major workplace factors associated with the condition are low job satisfaction and perceived poor workplace environment.¹¹

Classification of Neck Pain

There are many ways to categorize neck pain including duration (acute, <6 weeks; subacute, ≤3 months; chronic, >3 months), severity, etiology/structure, and type (ie, mechanical vs neuropathic). Among the various systems of categorization, duration is perhaps the best predictor of outcome. For a variety of different treatments, shorter duration has been found to be associated with a better prognosis than longstanding pain.¹²⁻¹⁴ The association between longer duration of pain and poorer prognosis is consistent with the findings in cohort studies that greater disease burden in general (eg, higher baseline pain scores and disability) predicts poorer outcomes for spinal pain.¹⁵⁻¹⁷

Neck pain can also be categorized by mechanisms as mechanical, neuropathic, or secondary to another cause (eg, referred pain from the heart or vascular pathology). Mechanical pain refers to pain originating in the spine or its supporting structures, such as ligaments and muscles. Common examples of mechanical pain include pain arising from the facet joints (eg, arthritis), diskogenic pain, and myofascial pain. Neuropathic pain refers to pain resulting

primarily from injury or disease involving the peripheral nervous system, which generally involves mechanical or chemical irritation of nerve roots. The most common examples of peripheral neuropathic pain are radicular symptoms from a herniated disk or osteophyte and spinal stenosis. Myelopathy, or symptoms arising from spinal cord pathology, is a form of central neuropathic pain. Mixed neuropathic-nociceptive pain states include postlaminectomy (failed neck surgery) syndrome and degenerated disks that result in a combination of mechanical pain from annular disruption and radicular symptoms from herniated nucleus pulposus (Figures 1-3).

Differentiating neuropathic from mechanical pain is probably the most important clinical distinction to make, as it affects treatment decisions at every level (eg, which medications, injections, or surgical procedure). There are several instruments available that have been found to distinguish neuropathic from nociceptive or mechanical pain, with 2 of the most common being the painDETECT questionnaire and the S-LANSS (Self-report Leeds Assessment of Neuropathic Symptoms




FIGURE 1. T2-weighted sagittal magnetic resonance image acquired slightly lateral to midline in a patient with unilateral radicular pain demonstrates a disk-osteophyte complex at C5-6 (arrow), which contributes to neural foraminal narrowing.



FIGURE 2. T2-weighted sagittal magnetic resonance image demonstrating multilevel disk bulging spanning levels C3-4 to C7-T1 causing central spinal stenosis in a patient with neuropathic pain extending into both arms. Ligamentum flavum hypertrophy at C5-6 (arrow A) and low-grade retrolisthesis of C4 on C5 (arrow B) contribute to central spinal stenosis. Note the absence of spinal cord signal hyperintensity, suggesting that there is no active cord edema.

and Signs) pain scale.^{18,19} For chronic low back pain, multiple studies have indicated a prevalence range of between 17% and 55% for predominantly neuropathic pain in a variety of cohorts, with a median of 41%.²⁰ No studies have examined the prevalence of neuropathic pain in a general neck pain population, but one study that aimed to validate S-LANSS and painDETECT in 152 individuals with cervical pain and a suspected nerve lesion found that 72% had definite or probable neuropathic pain, while another 18% had possible neuropathic pain according to the International Association for the Study of Pain Neuropathic Pain Special Interest Group grading system.²¹ Among 6 patients with whiplash, one-third had probable neuropathic pain and two-thirds possible neuropathic pain. Of note, the authors found that both instruments suffered from low sensitivities in this population.



FIGURE 3. T2-weighted sagittal magnetic resonance image in a patient with signs of myelopathy demonstrating a large central disk extrusion at C5-6. The signal hyperintensity within the spinal cord (arrow) indicates edema.

Natural Course of Neck Pain

Similar to back pain, most cases of acute (<6 weeks' duration) neck pain will resolve to a large extent within 2 months, but close to 50% of patients will continue to have some pain or frequent recurrences 1 year after occurrence.^{22,23} For acute pain, treatment appears to have little effect on persistence.²³ Factors that may be associated with poorer prognosis include female sex, older age, coexisting psychosocial pathology, and radicular symptoms (Table 1).²²⁻²⁶

A study by Gore et al²⁷ performed in patients with long-standing or recurrent neck pain found that individuals with more severe pain following an injury and those with symptoms or signs of cervical radiculopathy had a greater likelihood of persistent pain, although a formal statistical analysis was not performed for evaluation of radiculopathy. No association was found between the degree of radiographic degeneration and satisfaction with treatment results. A large retrospective, epidemiological study conducted in patients with radicular pain evaluated at Mayo Clinic found that although recurrence was

frequent (31.7%), at the mean follow-up of 5.9 years, 90.5% of patients experienced either no or only mild pain.²⁸ The finding that higher pain scores and radicular symptoms are associated with chronicity and poor outcome for neck pain is similar to what occurs with low back pain²⁹⁻³² and suggests that both subjective and objective factors play a role in prognosis.

The observation that most patients with cervical radiculopathy experience alleviation of symptoms with or without treatment is consistent with the results of small studies that revealed significant resorption in between 40% and 76% of cervical disk herniations.^{33,34} These statistics are similar to those noted for lumbar disk herniations.³⁵ Although acute neuropathic symptoms in spinal stenosis will stabilize or improve in over half of individuals, the anatomic derangements do not generally improve without treatment.^{36,37}

Cervical myelopathy involves pathology of the cervical spinal cord due to either trauma (spinal cord injury) or inflammation (myelitis), resulting in upper motor neuron signs. The natural course of nonsurgically treated myelopathy is highly variable. In a 1960s study that evaluated long-term follow-up in 28 patients treated nonoperatively, Lees and Turner³⁸ reported improvement in 17 patients, stable symptoms in 7, and progression in 4. Kadanka et al³⁹ conducted a randomized, 3-year study comparing surgical with nonsurgical treatment for mild to moderate spondylotic myelopathy. No differences were noted between treatment

TABLE 1. Factors Associated With the Development or Persistence of Neck Pain

Psychopathology
Low work satisfaction
Occupation/poor physical work environment
Female sex
Genetics
Concomitant back pain/other rheumatologic conditions
Poor coping skills
Catastrophization
Trauma/previous neck injury
Poor self-assessed health status
Sedentary lifestyle
Secondary gain
Smoking
Headache

groups, with 80% of patients in both groups exhibiting improvement or no clinical deterioration. Shimomura et al⁴⁰ also reported a 20% deterioration rate at a mean follow-up of 3 years. A prospective study by Sampath et al⁴¹ in 62 patients with cervical myelopathy found that equal proportions of medically and surgically treated patients (70%-75%) reported satisfaction with treatment, although the non-surgically treated patients experienced worsened neurologic symptoms and a decreased ability to perform activities of daily living. Some investigators have reported more dire outcomes for spondylotic myelopathy. Matsumoto et al⁴² reported that 10 of 27 patients treated conservatively over 6 months underwent surgery because of either neurologic deterioration or persistent disability. In another study by Sadasivan et al,⁴³ the authors reported deterioration in all 22 patients with cervical myelopathy, with 21 requiring surgery. In a consensus statement on the nonoperative treatment of spondylotic myelopathy, the authors concluded that between 20% and 62% of patients will have deterioration between 3-year and 6-year follow-up, with no patient or disease-specific factor being able to reliably predict progression of symptoms.⁴⁴

EVALUATION OF NECK PAIN

History

A comprehensive history can provide important clues regarding etiology and help differentiate primary neck pain from shoulder pain, thoracic outlet syndrome, brachial plexopathy, upper extremity pain, vascular pathology, and referred pain from thoracic viscera (eg heart, lungs). Patients with neuropathic pain typically describe their symptoms as shooting, electrical-like, stabbing, and/or burning, whereas mechanical pain is more often described as throbbing or aching.^{18,19} Neuropathic pain (eg, stenosis or herniated disk) is nearly always characterized by radiation into one or both upper extremities, usually in a single dermatomal or multidermatomal (eg, stenosis or a large or multilevel herniation) distribution. Because C7 and C6 are the most commonly affected nerve roots, radicular symptoms usually radiate into the middle or first 2 digits (eg, thumb and index finger), respectively.²⁸ Nonneuropathic pain arising from midlevel facet joints, disks

(eg, C5-6), or even muscles may also occasionally extend into the upper arm, but referral patterns tend to be nondermatomal and more variable.⁴⁵⁻⁴⁷ In pain stemming from the atlantoaxial, atlantooccipital, or upper facet joints or disks, radiation often extends into the occiput.⁴⁸

Associated signs and symptoms can often distinguish neuropathic from nonneuropathic pain. Neuropathic pain is frequently accompanied by numbness, paresthesias, or dysesthesias. Sensory symptoms are unusual in patients with nonneuropathic neck pain, and when they occur, they tend to be nondermatomal. The presence of confirmed neurologic symptoms in a patient with normal imaging results warrants a search for other sources of neuropathic pain, such as brachial plexopathy, or carpal or cubital tunnel syndrome.

Aggravating and alleviating factors can provide information relevant to the decision to pursue further work-up. Mechanical pain of any origin is classically associated with a low-level baseline pain that tends to worsen with activity, while neuropathic pain is associated with less predictable bouts of more intense exacerbations. Pain exacerbated when turning or bending the head ipsilateral to the source may indicate radicular or facetogenic pain, whereas pain worsened by contralateral turning of the head could suggest myofascial origin. Because the major cause of facet joint pain is arthritis, patients frequently report morning stiffness. Owing to a reduced spinal canal area, arm pain aggravated by neck extension is consistent with spinal stenosis; in contrast, pain worsened with forward flexion often signifies a diskogenic origin.

Cervical radiculopathy can often be distinguished from mechanical neck and shoulder pain by the abduction relief sign, in which abduction of the ipsilateral arm over the head alleviates symptoms.⁴⁹ This maneuver can distinguish radicular from certain types of shoulder pain, which may be worsened by shoulder abduction. One condition that is often mistaken for cervical radicular pain is thoracic outlet syndrome, which may be neurogenic (which comprises about 95% cases), arterial, and/or venous in origin. Thoracic outlet syndrome is classically unilateral, affects women more frequently than men, and peaks in prevalence in the fourth decade of life. In about half the cases, it is preceded by either trauma or

repetitive stress. Imaging and Doppler analysis are most helpful for diagnosing vascular thoracic outlet syndrome but have low sensitivity for the neurogenic type. Several tests have been advocated to identify thoracic outlet syndrome including the elevated arm stress test, Adson test, and tenderness to palpation at the scalene triangle or insertion of the pectoralis minor, although none have high specificity.⁵⁰ Because neck pain is typically alleviated by rest and recumbency, severe unrelenting pain not affected by rest or position warrants consideration of “red flags” such as malignant neoplasms, primary neurologic conditions, and infection (Table 2, Figure 4).

Occasionally, the inciting event can facilitate identification of a pain generator. The most common precipitating event for neck pain is whiplash injury, which occurs when the neck and head continue to lurch forward after the trunk has ceased to move, resulting in shearing stress that involves the disks and facet joints that connect adjacent vertebrae. Although Bogduk and Yoganandan⁵¹ reported that videoradiographic studies performed with and without headrests

in cadavers in the 1970s indicated that rear-end collisions were most frequently associated with injuries to the intervertebral disks (90%), anterior spinal ligaments (80%), and facet joints (40%), more recent⁵² and methodologically sound⁵³ studies have found no consistent relationship between pain and imaging abnormalities following motor vehicle collisions. In clinical studies performed by the Bogduk group using response to “double-blocks” as the reference standard, between 30% and 60% of patients with whiplash injury have predominantly facet joint pain.⁵⁴⁻⁵⁶ For nontraumatic facetogenic and discogenic pain, the onset tends to be insidious because of the progressive strain on these structures from repetitive, low-level stress. For cervical radicular pain, particularly in younger individuals with robust disks, patients will sometimes report a specific antecedent event.

Physical Examination

The physical examination is often used to confirm a historical finding, screen patients for serious or treatable pathology, and inform advanced imaging or further diagnostic work-

TABLE 2. What Not to Miss: “Red Flags” Associated With Neck Pain

Red flag	Potential conditions	Associated signs and symptoms
Trauma (eg, fall, motor vehicle accident, whiplash injury)	Vertebral fractures, spinal cord injury/syrinx, ligamentous disruption	Loss of or alternating consciousness, cognitive deficits, traumatic brain injury, headaches, neurologic symptoms
Rheumatoid arthritis, Down syndrome, spondyloarthropathy	Atlantoaxial subluxation	Easy fatigability, gait abnormalities, limited neck mobility, torticollis, clumsiness, spasticity, sensory deficits, upper motor neuron signs
Constitutional symptoms	Metastases, infectious process, systemic rheumatologic disease	Weight loss, unexplained fevers, anorexia, family or personal history of malignant neoplasm, diffuse joint pain and stiffness, abnormal laboratory test results
Infectious symptoms	Epidural abscess, spondylodiskitis, meningitis	Fever, neck stiffness, photophobia, elevated white blood cell count
Upper motor neuron lesion	Spinal cord compression, demyelinating disease	Hoffmann sign, hyperreflexia, Babinski sign, spasticity, incontinence, sexual dysfunction
Age <20 y	Congenital abnormalities (cervical spina bifida, Scheuermann disease), conditions associated with substance abuse such as infection	Congenital anomalies: birthmarks, overlying skin tags, patches of hair, family history, systemic disease (eg, diabetes, epilepsy for spina bifida) Substance abuse: male sex, poor work or school performance, depression or other psychiatric morbidity
Concurrent chest pain, diaphoresis, or shortness of breath	Myocardial ischemia or infarction	Nausea, extension of pain into the left arm (especially medial upper arm)
Age >50 y	Metastases, vertebral fracture, carotid or vertebral artery dissection/bleeding	Family or personal history of malignant neoplasm, previous trauma Arterial dissection: tearing sensation, headache, visual loss, or other neurologic sequelae



FIGURE 4. T1-weighted sagittal magnetic resonance image in a patient with a known primary malignant neoplasm demonstrating hypointense lesions within the C3 (arrow A) and T3 (arrow B) vertebral bodies. At T3, there is an accompanying compression fracture with loss of vertebral body height. Breast, lung, prostate, renal cell, and gastrointestinal tract cancers, lymphoma, and melanoma are the primary malignant neoplasms most likely to metastasize to the vertebral bodies and should be considered in the differential diagnosis of a vertebral body infiltrative lesion in patients older than 40 years of age.

up but is rarely pathognomonic. Gait abnormalities, which can herald spinal cord (eg, myelopathy or syrinx) or brain injury, and major traumatic or developmental abnormalities should be noted. For example, doughy lipomata may indicate spina bifida or spinal cord abnormalities, and a prominent, palpable vertebral body can signify spondylolisthesis.

General appearance should be observed to identify facial expressions and behaviors indicative of pain. Patients who report severe pain in the absence of pain-related behaviors should be further evaluated for signs of nonorganic pathology. Abnormal lateral or forward flexion, or rotation, may indicate torticollis. Muscle atrophy, or winging or drooping of the shoulder, may be observed with radiculopathy, brachial plexopathy, or nerve entrapment.

True neurologic weakness should be distinguished from pain-induced weakness. In individuals with nerve injury, muscle wasting or asymmetric reflexes may be present, although 10% of asymptomatic individuals may have absent or asymmetric reflexes. In patients with poor effort or suspected malingering, reflexes may be the most (or only) objective examination tool. Signs of upper motor neuron lesions must be vigorously investigated.

Range of motion may be limited in all types of mechanical neck pain, but specific exacerbating movements may provide clues to the origin. For example, reproducible arm pain with neck flexion toward the affected side may indicate foraminal stenosis and/or radiculopathy. In one study conducted in whiplash patients, no difference in facet block responders and nonresponders was found for range of motion in any direction.⁵⁷ Provocative maneuvers may be more helpful in identifying potential sources of neuropathic pain. For cervical radiculopathy, the Spurling shoulder abduction and neck distraction tests have moderate sensitivity (approximately 50%) but high specificity (>80%).^{58,59} For cervical myelopathy, the Hoffmann sign has been reported to have moderate sensitivity and specificity.^{59,60} For facetogenic pain, one study found that paraspinal tenderness was weakly correlated with positive treatment response⁶¹ (Tables 3 and 4).

Diagnostic Work-up

In patients with suspected structural abnormalities (eg, scoliosis, spondylolisthesis, fractures), plain radiographs are generally sufficient. Magnetic resonance imaging (MRI) is the most sensitive test for detecting soft-tissue (eg, disk) abnormalities but is characterized by a high rate of abnormalities in asymptomatic individuals. The rates of abnormalities in people without symptoms varies from around 60% in individuals in their 40s to more than 80% in individuals older than 60 years, with the most common abnormalities being decreased signal intensity and disk protrusions.^{62,63} Therefore, MRI is recommended to rule out red flags, in patients with serious or progressive neurologic deficits, and when referring patients for procedural interventions (eg, surgery); for individuals with persistent pain that does not

TABLE 3. Accuracy of Physical Examination Tests for Neck Pain

Test	Description	Diagnosis	Accuracy ^a
Spurling	Lateral flexion and rotation to the affected side with axial compression of the head reproduces radicular pain	Cervical radiculopathy	40%-60% Sensitivity, 85%-95% specificity; moderate to substantial reliability
Shoulder abduction	Relief of ipsilateral cervical radicular symptoms with abduction of symptomatic arm (eg, placing it on head)	Cervical radiculopathy	40%-50% Sensitivity, 80%-90% specificity; fair to moderate reliability
Neck distraction	Relief of radicular symptoms when examiner grasps patient's head under occiput and chin and lifts, applying axial traction	Cervical radiculopathy	40%-50% Sensitivity, 90% specificity; moderate reliability
Valsalva	Reproduction of radicular pain with forced expiratory effort with mouth and nose closed	Cervical radiculopathy	Low sensitivity (22%), high specificity (94%)
Upper limb tension	Reproduction of radicular pain with scapular depression; shoulder abduction; forearm supination, wrist and finger extension; shoulder external rotation; elbow extension; contralateral followed by ipsilateral cervical lateral flexion	Cervical radiculopathy	70%-90% Sensitivity, 15%-30% specificity
Lhermitte sign	Electrical-like sensations down spine or arms with passive flexion of neck	Cervical myelopathy	<20% Sensitivity, >90% specificity
Hoffmann sign	Flexion-adduction of thumb and index finger elicited with snapping flexion of middle or fourth finger distal phalanx	Cervical myelopathy	50%-80% Sensitivity, 78% specificity
Babinski sign	Stimulation of the sole of the foot elicits dorsiflexion of hallux and sometimes dorsiflexion and abduction of other toes	Cervical myelopathy	10%-75% Sensitivity, >90% specificity
Hyperreflexia	Overreactive or overresponsive deep tendon reflexes	Cervical myelopathy above level of muscle reflex innervation	>65% Sensitivity, high specificity
Clonus	>2 Repetitive beats during wrist or ankle dorsiflexion movements	Cervical myelopathy	<50% Sensitivity
Jackson compression	Downward pressure on head with lateral flexion. Localized pain may indicate facet joint pain; arm pain may indicate radiculopathy	Cervical radiculopathy/myelopathy or facet joint pain	Not validated for facet joint pain. Low sensitivity, high specificity for myelopathy
Paraspinal tenderness	Paraspinal > midline pain with palpation	Cervical facet joint pain	Weak evidence for predicting a positive response to treatment

^aRanges of accuracy are given when multiple studies were available. Data from *Eur Spine J*,⁵⁸ *Pain Physician*,⁵⁹ and *Phys Ther*.⁶⁰

respond to conservative treatment, radiologic evaluations can be considered.

Electrodiagnostic testing can be considered in patients with equivocal symptoms or imaging findings and to rule out peripheral neuropathy. The American Association of Electrodiagnostic Medicine reported 50% to 71% sensitivity in diagnosing cervical radiculopathy,⁶⁴ but a later study by Ashkan et al⁶⁵ found that compared with neurophysiologic studies, MRI was associated with a higher sensitivity (93% vs 42%) and negative predictive value (25% vs 7%) based on operative findings. Selective nerve root blocks have been used to identify painful nerve root(s)

and have been reported to improve surgical outcomes, but randomized trials are lacking (see "Injections" section).⁶⁶⁻⁶⁸

TREATMENT

Conservative Therapy

Similar to back pain, cervical and scapular stretching and strengthening exercises have been found to provide intermediate-term relief for mechanical neck pain.⁶⁹⁻⁷¹ In one large randomized study of 206 patients with acute cervical radiculopathy, both physical therapy accompanied by home exercises and the use of a hard

TABLE 4. Signs and Symptoms of Cervical Radiculopathy

Affected nerve root (frequency) ^a	Pain location	Sensory deficits	Muscle weakness	Reflex abnormalities
C4 (<10%)	Upper-mid neck	Capelike distribution, shoulder	None	None
C5 (10%)	Neck, shoulder, interscapular region, anterior arm	Lateral aspect of shoulder and arm	Shoulder abduction and external rotation, elbow flexion	Deltoid, biceps, and brachioradialis
C6 (20%-25%)	Neck, shoulder, interscapular region, lateral forearm, first and second digits	Lateral aspect of forearm and hand, first and second digits	Elbow flexion, shoulder external rotation, abduction and protraction, forearm supination and pronation, wrist extension	Biceps, brachioradialis
C7 (45%-60%)	Lower neck, shoulder, interscapular region, extensor surface of forearm, chest, third digit	Third digit, sometimes parts of first 4 digits	Elbow and finger extension, forearm pronation	Triceps
C8 (10%)	Lower neck, medial forearm and hand	Distal medial forearm to medial hand and fourth and fifth digits	Wrist flexion, finger and thumb abduction, adduction, extension and flexion	Finger flexors

^aPercentage data in part from *Brain*.²⁸

cervical collar produced greater reductions in neck pain and disability over a 6-week period than a “wait and see” approach.⁷² However, systematic reviews have concluded that cervical collars are no more effective than sham interventions for neck pain.⁷³ For complementary and alternative medical treatments, the strongest evidence supports a modest effect for spinal manipulation compared with no treatment or other noninterventional treatments. With regard to other complementary and alternative treatments, although they have generally been found to be superior to no treatment, the evidence that they are superior to sham treatments or other treatments is weak, negative, or conflicting (Table 5).

Few high-quality studies have evaluated pharmacotherapy for neck pain. Systemic nonsteroidal anti-inflammatory drugs (NSAIDs) have been found to be beneficial for spinal pain in general⁸² but have not been formally studied in neck pain. Although NSAIDs are more efficacious than acetaminophen, the American College of Rheumatologists recommends acetaminophen as a first-line treatment, even for arthritis, because of its more favorable adverse effect profile.⁸³ In patients who present with predominantly mechanical neck pain, topical NSAIDs have proven efficacy.⁸⁴ In one randomized trial that compared spinal manipulation, home

exercise and advice, and pharmacotherapy with NSAIDs or acetaminophen in acute and subacute neck pain, the manipulation and exercise groups fared better than medicinal treatment through 12-month follow-up.⁸¹

In 2 large (n=1405) randomized controlled trials evaluating the muscle relaxant cyclobenzaprine for acute neck (more than one-third of the patients) or low back pain associated with muscle spasm, the authors found both intermediate-dose (15 mg/d) and high-dose (30 mg/d) therapy to be more effective than placebo but no difference between low doses (7.5 mg/d) and placebo.⁸⁵ A double-blind crossover study comparing the stand-alone anti-inflammatory drug benorylate to benorylate plus the muscle relaxant chlormezanone found no benefit of add-on therapy for low back or joint pain but significantly better pain relief and sleep quality in patients with neck pain.⁸⁶ Muscle relaxants tend to be more effective for acute than chronic pain.⁸⁷ In light of their abuse potential and lack of greater efficacy compared with other muscle relaxants, many experts believe benzodiazepines should be prescribed only when other muscle relaxants have proven ineffective and with clearly defined goals, time frames, and appropriate surveillance.⁸⁸

TABLE 5. Alternative and Complementary Medicine Treatments for Neck Pain

Treatment	Description	Evidence
Spinal manipulation	Manual therapy designed to maximize painless movement, reduce muscle tightness, improve joint mobility, and correct alignment problems	Superior to no treatment or sham treatment in the short term. Weak evidence for intermediate-term benefit and for superiority over pharmacotherapy and other alternative therapies
Acupuncture	Inserting needles into the skin at various anatomic locations to reduce pain or induce anesthesia. Needles may be manipulated manually or through electrical stimulation	Weak evidence that acupuncture is superior to no treatment in the short term. Strong evidence that acupuncture is not better than sham acupuncture or other treatments
Massage therapy	The manipulation of muscle and connective tissue to enhance function and promote relaxation and well-being	Superior to no treatment or sham treatment but not more effective than other active treatments in the short and intermediate term. No evidence for improved function
Exercise therapy	Active or passive physical exercises designed to strengthen or stabilize the spine that may reduce pain, prevent injuries, and improve posture and body mechanics	Strong evidence for intermediate-term relief for nonspecific neck pain and whiplash-type injuries. Conflicting evidence for improvement of disability. No clear evidence supporting one technique over another or that exercise can prevent the development of neck pain
Traction	Procedures designed to relieve pressure on the spine	There is low-quality evidence that traction is not superior to placebo treatments for neck pain with or without radiculopathy
Soft cervical collar	Orthopedic device used to immobilize the neck and support the head and neck, often after injury	There is low-quality evidence that a cervical collar is no more effective than physical therapy or other active therapies for cervical radiculopathy and whiplash
Electrotherapy	The use of electrical energy as a medical treatment to relieve pain, usually by interfering with nerve conduction	There is low-quality evidence that various forms of electrotherapy (eg, transcutaneous electrical nerve stimulation, pulsed electromagnetic field therapy) are better than placebo but not other treatments
Yoga	A series of physical, mental, and spiritual exercises designed to achieve a peaceful state of mind, improve conditioning, and attain self-actualization	There is weak evidence that yoga is more effective than home-based exercise treatment

Data from references 69 through 81.

Injections

The evidence supporting trigger point injections to treat myofascial pain is mixed. Part of the difficulty in evaluating clinical trials for trigger point injections is that the injection of any substance (or even dry needling) into taut bands of muscle may relieve pain, which makes it difficult to perform true placebo-controlled trials. In a systematic review by Scott et al⁸⁹ evaluating trigger point injections for chronic pain, the authors found no clear evidence for either benefit or ineffectiveness. With regard to the type of procedure, there is limited evidence that injections may be more effective and less painful than dry needling.^{89,90} For botulinum toxin, there is mixed evidence for superiority over trigger point injections performed with saline or local anesthetic. A Cochrane review identified 4 studies that met

inclusion criteria, 3 of which focused on myofascial pain in the neck and/or shoulder region.⁹¹ Although all 3 studies favored botulinum toxin, in only 1 study did the results reach statistical significance.⁹²⁻⁹⁴ A more recent study of 114 previous responders that used an enriched protocol design found modest benefit in some but not most outcome measures for some variables that persisted through 26 weeks.⁹⁵ In 2 controlled studies performed in patients with low back pain, the results were mixed regarding the effectiveness of botulinum toxin.^{96,97} The authors found there was inconclusive evidence to support the use of botulinum toxin injections for myofascial pain syndrome. In a non-systematic review by Jabbari and Machado,⁹⁸ the authors concluded there was level A evidence for the use of botulinum toxin injections to treat cervical dystonia. One caveat that is important to

heed when interpreting studies on trigger point injections is that they are widely acknowledged to be more effective when used in patients in whom discrete, taut bands of muscle can be palpated (ie, trigger points) than in individuals with more diffuse symptoms.⁴⁷ In some primary studies, however, the methodology used to identify trigger points was unclear.

For cervical radiculopathy, the results of clinical trials evaluating epidural corticosteroid injections have been mixed. A small (n=40) randomized study found no significant differences at 3-week follow-up between transforaminal corticosteroids plus local anesthetic and transforaminal local anesthetic.⁹⁹ This is consistent with 3 randomized, double-blind studies by the same group that compared epidural corticosteroid plus local anesthetic to epidural local anesthetic alone in a variety of conditions (herniated disc, spinal stenosis, and failed neck surgery syndrome) and found no differences between treatment groups, with both groups experiencing improvement.¹⁰⁰⁻¹⁰² In a randomized, placebo-controlled, nonblinded study that compared a series of epidural corticosteroid and local anesthetic injections to intramuscular injections, Stav et al¹⁰³ reported significant benefit lasting up to 1 year. In a large, multicenter comparative effectiveness study, Cohen et al¹⁰⁴ found that combination treatment with a series of epidural corticosteroid injections plus conservative treatment with adjuncts and physical therapy was superior to either treatment alone. Of note, a systematic review and meta-analysis concluded that epidural local anesthetic and/or saline constituted an efficacious treatment intermediate in efficacy between epidural corticosteroids and a true intramuscular placebo injection.¹⁰⁵

Cervical facet joint pain is estimated to account for between 40% and 60% of nonneuropathic neck pain based on controlled blocks.^{55,106} The evidence for medial branch (facet joint nerve) radiofrequency ablation of cervical facet joint pain is weakly positive. In one small, placebo-controlled study performed in 24 meticulously selected patients with whiplash injury, the treatment group fared better than the sham group for pain relief and functional improvement, with the mean duration of benefit lasting about 9 months.¹⁰⁷ In a smaller study in 12 patients with

cervicogenic headache that performed empirical radiofrequency denervation without diagnostic injections, 4 of 6 persons in the treatment group experienced success at 3 months, which favorably compared with 2 of 6 in the treatment group.¹⁰⁸ Although small uncontrolled studies have reported benefit with intra-articular corticosteroid injections,¹⁰⁹ the only placebo-controlled study reported no differences between the corticosteroid and local anesthetic control injections at 6-month follow-up.¹¹⁰

Injections in the form of selective nerve root blocks (SNRBs) have also been advocated as a tool to identify symptomatic spinal levels and select patients for surgery.⁶⁸ There is a strong correlation between the results of SNRB and single-level MRI pathology,¹¹¹ but the correlation between SNRBs, MRI findings, and neurologic examination results in individuals with multilevel pathology is much lower.⁶⁶ Although uncontrolled studies have found good surgical outcomes in patients who experience pain relief after diagnostic injections,^{66,67,111} there have been no randomized studies evaluating their ability to improve treatment results.⁶⁸ One recent review concluded that adding SNRB to diagnostic work-ups in patients with lumbar radiculopathy being considered for surgery was not cost-effective.¹¹²

Surgery

Few randomized studies have evaluated surgical treatment for neck pain, and none have done so for mechanical pain. In a randomized study comparing anterior decompression and fusion operations, physical therapy, and hard collar immobilization in 81 patients with cervical radiculopathy, Persson et al^{113,114} found greater reductions in pain (29% for surgery, 19% for physical therapy, and 4% for cervical collar) and improvements in muscle strength and sensory loss in the surgical group than in the other treatment groups. Yet at 1-year follow-up, the differences favoring surgery were for the most part no longer statistically significant. A more recent randomized study that compared surgery and physical therapy to physical therapy alone for cervical radiculopathy found that surgery was associated with superior outcomes at 1 year, but by 2 years, the differences between groups were no longer statistically significant.¹¹⁵ In a clinical trial

performed in 120 patients with neck and/or arm pain secondary to a single, small contained disk herniation, plasma disk decompression was found to be superior to conservative treatment for pain and function throughout the 1-year follow-up.¹¹⁶ Of note, conservative treatment had already failed in all patients.

In the only randomized study evaluating surgery for cervical myelopathy, Kadanka et al¹¹⁷ compared operative therapy to conservative care consisting of immobilization with a soft collar, NSAIDs, and intermittent bed rest. Sixty-eight patients were randomized by coin flip, with discrepancies noted for some baseline variables. Overall, through the 10-year follow-up period, no significant differences were found for major outcome variables between treatment groups.^{117,118} A subgroup analysis found that patients who were younger and had greater baseline disease burden and small spinal canal areas tended to fare better with surgery than those who were older and had greater function and transverse spinal canal diameter.¹¹⁹

There are no randomized controlled trials comparing surgical to nonsurgical therapies for mechanical neck pain associated with common degenerative changes, but extrapolated studies in the lumbar spine suggest that less than one-third of patients will experience clinically meaningful pain relief or functional improvement, with the results diminishing over time.^{120,121} The results of systematic reviews comparing cervical disk replacement to anterior decompression and fusion operations are conflicting as to whether the former is associated with better outcomes for single-level spondylosis.^{122,123} One study evaluating outcome predictors for anterior cervical decompression and fusion found that good functional capacity, male sex, and nonsmoking status were associated with successful long-term treatment results.¹²⁴

FUTURE DIRECTIONS

Compared with other leading causes of pain and disability, relatively few randomized controlled trials exist to guide treatment of neck pain, and the guidelines for neck pain are often extrapolated from those for other conditions. Clinical trials designed to determine efficacy and comparative effectiveness are needed for all types of treatments but particularly

adjuvants for neuropathic pain and surgery for mechanical pain.

The use of biological therapies, including stem cell therapy, and nerve growth factor and cytokine inhibitors have been or are currently being studied for other chronic pain conditions such as low back pain but have yet to be critically evaluated for neck pain. Future research should be expanded to determine their efficacy for spinal pain in general or neck pain in particular.

The persistence of neck pain after whiplash and other types of injuries poses substantial physical, psychological, and economic consequences for patients and society. There is currently a very poor relationship between symptoms and imaging abnormalities in injured patients who continue to experience neck pain.⁵² Finding ways to identify those individuals at increased risk for development of persistent pain, and preventing it, represents an important challenge to the medical community.

CONCLUSION

Neck pain is one of the leading causes of disability in the world, yet the amount of research devoted to treatment is relatively low in comparison to the other leading causes. For acute neck pain, most cases will resolve spontaneously over a period of weeks to months, but a substantial proportion of individuals will be left with residual or recurrent symptoms. Treatment appears to have little effect on the course of acute neck pain. History and physical examination may provide important clues as to whether the pain is neuropathic or mechanical and are critical in determining who might benefit from advanced imaging or further diagnostic work-up. In patients with whiplash injuries, there is a poor correlation between pain and imaging results. Clinical trials have found that exercise may be beneficial, and for acute pain, muscle relaxants are effective. In individuals with chronic pain, there is conflicting evidence supporting epidural corticosteroid injections in patients with radiculopathy and spinal stenosis and weak evidence in favor of facet joint radiofrequency denervation for spinal arthritis.

ACKNOWLEDGMENTS

The opinions or assertions contained herein are the private views of the author and are

not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Abbreviations and Acronyms: MRI = magnetic resonance imaging; NSAID = nonsteroidal anti-inflammatory drug; SNRB = selective nerve root block

Grant Support: This work was funded in part by a Congressional grant from the Center for Rehabilitation Sciences Research. The role of the funding source was only provisions to pay research personnel.

Potential Competing Interests: Dr Cohen serves as a consultant for Regenesi Biomedical and Semnur Pharmaceuticals, Inc, and is on the advisory board of Kimberly Clark Health Care.

Correspondence: Address to Steven P. Cohen, MD, 550 N Broadway, Ste 301, Baltimore, MD 21029 (scohen40@jhmi.edu). Individual reprints of this article and a bound reprint of the entire Symposium on Pain Medicine may be available for purchase from our website www.mayoclinicproceedings.org.

The Symposium on Pain Medicine will continue in an upcoming issue.

REFERENCES

1. US Burden of Disease Collaborators. The state of US health, 1990-2010: burden of diseases, injuries, and risk factors. *JAMA*. 2013;310(6):591-608.
2. Fejer R, Kyvik KO, Hartvigsen J. The prevalence of neck pain in the world population: a systematic critical review of the literature. *Eur Spine J*. 2006;15(6):834-848.
3. Hogg-Johnson S, van der Velde G, Carroll LJ, et al. Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. The burden and determinants of neck pain in the general population: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)*. 2008;33(4, suppl):S39-S51.
4. Binder AI. Neck pain. *Clin Evid (online)*. 2008;pii:1103.
5. Fernández-de-las-Peñas C, Hernández-Barrera V, Alonso-Blanco C, et al. Prevalence of neck and low back pain in community-dwelling adults in Spain: a population-based national study. *Spine (Phila Pa 1976)*. 2011;36(3):E213-E219.
6. Strine TW, Hootman JM. US national prevalence and correlates of low back and neck pain among adults. *Arthritis Rheum*. 2007;57(4):656-665.
7. Son KM, Cho NH, Lim SH, Kim HA. Prevalence and risk factor of neck pain in elderly Korean community residents. *J Korean Med Sci*. 2013;28(5):680-686.
8. Nilsen TI, Holtermann A, Mork PJ. Physical exercise, body mass index, and risk of chronic pain in the low back and neck/shoulders: longitudinal data from the Nord-Trøndelag Health Study. *Am J Epidemiol*. 2011;174(3):267-273.
9. Käänä S, Laaksonen M, Rahkonen O, Lahelma E, Leino-Arjas P. Risk factors of chronic neck pain: a prospective study among middle-aged employees. *Eur J Pain*. 2012;16(6):911-920.
10. Vincent HK, Adams MC, Vincent KR, Hurley RW. Musculoskeletal pain, fear avoidance behaviors, and functional decline in obesity: potential interventions to manage pain and maintain function. *Reg Anesth Pain Med*. 2013;38(6):481-489.
11. Côté P, van der Velde G, Cassidy JD, et al. The burden and determinants of neck pain in workers: results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *J Manipulative Physiol Ther*. 2009;32(2, suppl):S70-S86.
12. May S, Gardiner E, Young S, Kluber-Moffett J. Predictor variables for a positive long-term functional outcome in patients with acute and chronic neck and back pain treated with a McKenzie approach: a secondary analysis. *J Man Manip Ther*. 2008;16(3):155-160.
13. Royuela A, Kovacs FM, Campillo C, Casamitjana M, Muriel A, Abreira V. Predicting outcomes of neuroreflexotherapy in patients with subacute or chronic neck or low back pain. *Spine J*. 2014;14(8):1588-1600.
14. Peterson C, Bolton J, Humphreys BK. Predictors of outcome in neck pain patients undergoing chiropractic care: comparison of acute and chronic patients. *Chiropr Man Therap*. 2012;20(1):27.
15. Enthoven P, Skargren E, Carstensen J, Oberg B. Predictive factors for 1-year and 5-year outcome for disability in a working population of patients with low back pain treated in primary care. *Pain*. 2006;122(1-2):137-144.
16. Wilkens P, Scheel IB, Grundnes O, Hellum C, Storheim K. Prognostic factors of prolonged disability in patients with chronic low back pain and lumbar degeneration in primary care: a cohort study. *Spine (Phila Pa 1976)*. 2013;38(1):65-74.
17. Kleinstueck FS, Fekete T, Jeszenszky D, et al. The outcome of decompression surgery for lumbar herniated disc is influenced by the level of concomitant preoperative low back pain. *Eur Spine J*. 2011;20(7):1166-1173.
18. Freynhagen R, Baron R, Gockel U, Tölle TR. painDETECT: a new screening questionnaire to identify neuropathic components in patients with back pain. *Curr Med Res Opin*. 2006;22(10):1911-1920.
19. Bennett M. The LANS Pain Scale: the Leeds Assessment of Neuropathic Symptoms and Signs. *Pain*. 2001;92(1-2):147-157.
20. Cohen SP, Bicket MC, Jamison D, Wilkinson I, Rathmell JP. Epidural steroids: a comprehensive, evidence-based review. *Reg Anesth Pain Med*. 2013;38(3):175-200.
21. Tampin B, Briffa NK, Goucke R, Slater H. Identification of neuropathic pain in patients with neck/upper limb pain: application of a grading system and screening tools. *Pain*. 2013;154(12):2813-2822.
22. Vasseljen O, Woodhouse A, Bjørggaard JH, Leivseth L. Natural course of acute neck and low back pain in the general population: the HUNT study. *Pain*. 2013;154(8):1237-1244.
23. Vos CJ, Verhagen AP, Passchier J, Koes BW. Clinical course and prognostic factors in acute neck pain: an inception cohort study in general practice. *Pain Med*. 2008;9(5):572-580.
24. Pemo G, Mortimer M, Wiktorin C, Tomqvist EW, Vingård E; Musculoskeletal Intervention Center-Norrålle Study Group. Neck/shoulder disorders in a general population: natural course and influence of physical exercise; a 5-year follow-up. *Spine (Phila Pa 1976)*. 2005;30(13):E363-E368.
25. Christensen JO, Knardahl S. Time-course of occupational psychological and social factors as predictors of new-onset and persistent neck pain: a three-wave prospective study over 4 years. *Pain*. 2014;155(7):1262-1271.
26. Carstensen TB. The influence of psychosocial factors on recovery following acute whiplash trauma. *Dan Med J*. 2012;59(12):B4560.
27. Gore DR, Sepic SB, Gardner GM, Murray MP. Neck pain: a long-term follow-up of 205 patients. *Spine (Phila Pa 1976)*. 1987;12(1):1-5.
28. Radhakrishnan K, Litchy WJ, O'Fallon WM, Kurland LT. Epidemiology of cervical radiculopathy: a population-based study from Rochester, Minnesota, 1976 through 1990. *Brain*. 1994;117(pt 2):325-335.

29. Hsu E, Atanelov L, Plunkett AR, Chai N, Chen Y, Cohen SP. Epidural lysis of adhesions for failed back surgery and spinal stenosis: factors associated with treatment outcome. *Anesth Analg*. 2014;118(1):215-224.
30. Costa Lda C, Maher CG, McAuley JH, et al. Prognosis for patients with chronic low back pain: inception cohort study. *BMJ*. 2009;339:b3829.
31. Cherkin DC, Deyo RA, Street JH, Barlow W. Predicting poor outcomes for back pain seen in primary care using patients' own criteria. *Spine (Phila Pa 1976)*. 1996;21(24):2900-2907.
32. Thomas E, Silman AJ, Croft PR, Papageorgiou AC, Jayson MI, Macfarlane GJ. Predicting who develops chronic low back pain in primary care: a prospective study. *BMJ*. 1999;318(7199):1662-1667.
33. Maigne JY, Deligne L. Computed tomographic follow-up study of 21 cases of nonoperatively treated cervical intervertebral soft disc herniation. *Spine (Phila Pa 1976)*. 1994;19(2):189-191.
34. Mochida K, Komori H, Okawa A, Muneta T, Haro H, Shinomiya K. Regression of cervical disc herniation observed on magnetic resonance images. *Spine (Phila Pa 1976)*. 1998;23(9):990-995.
35. Saal JA. Natural history and nonoperative treatment of lumbar disc herniation. *Spine (Phila Pa 1976)*. 1996;21(24, suppl):25-95.
36. Minamide A, Yoshida M, Maio K. The natural clinical course of lumbar spinal stenosis: a longitudinal cohort study over a minimum of 10 years. *J Orthop Sci*. 2013;18(5):693-698.
37. Micankova Adamova B, Vohanka S, Dusek L, Jarkovsky J, Bednarik J. Prediction of long-term clinical outcome in patients with lumbar spinal stenosis. *Eur Spine J*. 2012;21(12):2611-2619.
38. Lees F, Turner JW. Natural history and prognosis of cervical spondylosis. *Br Med J*. 1963;2(5373):1607-1610.
39. Kadanka Z, Mareš M, Bednarik J, et al. Approaches to spondylotic cervical myelopathy: conservative versus surgical results in a 3-year follow-up study. *Spine (Phila Pa 1976)*. 2002;27(20):2205-2210.
40. Shimomura T, Sumi M, Nishida K, et al. Prognostic factors for deterioration of patients with cervical spondylotic myelopathy after nonsurgical treatment. *Spine (Phila Pa 1976)*. 2007;32(22):2474-2479.
41. Sampath P, Bendebba M, Davis JD, Ducker TB. Outcome of patients treated for cervical myelopathy: a prospective, multicenter study with independent clinical review. *Spine (Phila Pa 1976)*. 2000;25(6):670-676.
42. Matsumoto M, Chiba K, Ishikawa M, Maruiwa H, Fujimura Y, Toyama Y. Relationships between outcomes of conservative treatment and magnetic resonance imaging findings in patients with mild cervical myelopathy caused by soft disc herniations. *Spine (Phila Pa 1976)*. 2001;26(14):1592-1598.
43. Sadasivan KK, Reddy RP, Albright JA. The natural history of cervical spondylotic myelopathy. *Yale J Biol Med*. 1993;66(3):235-242.
44. Fehlings MG, Wilson JR, Yoon ST, Rhee JM, Shamji MF, Lawrence BD. Symptomatic progression of cervical myelopathy and the role of nonsurgical management: a consensus statement. *Spine (Phila Pa 1976)*. 2013;38(22, suppl 1):S19-S20.
45. Fukui S, Ohseto K, Shiotani M, et al. Referred pain distribution of the cervical zygapophyseal joints and cervical dorsal rami. *Pain*. 1996;68(1):79-83.
46. April C, Dwyer A, Bogduk N. Cervical zygapophyseal joint pain patterns, II: A clinical evaluation. *Spine (Phila Pa 1976)*. 1990;15(6):458-461.
47. Simons DG, Travell JG, Simons LS. In: . *Travell and Simons' Myofascial Pain and Dysfunction: The Trigger Point Manual*. 2nd ed., Vol. 1. Baltimore, MD: Williams & Wilkins; 1999: 11-93.
48. Dreyfuss P, Michaelsen M, Fletcher D. Atlanto-occipital and lateral atlanto-axial joint pain patterns. *Spine (Phila Pa 1976)*. 1994;19(10):1125-1131.
49. Fast A, Parikh S, Marin EL. The shoulder abduction relief sign in cervical radiculopathy. *Arch Phys Med Rehabil*. 1989;70(5):402-403.
50. Freischlag J, Orion K. Understanding thoracic outlet syndrome. *Scientifica (Cairo)*. 2014;2014:248163.
51. Bogduk N, Yoganandan Y. Biomechanics of the cervical spine Part 3: minor injuries. *Clin Biomech (Bristol, Avon)*. 2001;16(4):267-275.
52. Matsumoto M, Okada E, Ichihara D, et al. Prospective ten-year follow-up study comparing patients with whiplash-associated disorders and asymptomatic subjects using magnetic resonance imaging. *Spine (Phila Pa 1976)*. 2010;35(18):1684-1690.
53. Pettersson K, Hildingsson C, Toolanen G, Fagerlund M, Björnebrink J. MRI and neurology in acute whiplash trauma: no correlation in prospective examination of 39 cases. *Acta Orthop Scand*. 1994;65(5):525-528.
54. Bamsley L, Lord S, Bogduk N. Comparative local anaesthetic blocks in the diagnosis of cervical zygapophysial joint pain. *Pain*. 1993;55(1):99-106.
55. Lord SM, Bamsley L, Wallis BJ, Bogduk N. Chronic cervical zygapophysial joint pain after whiplash: a placebo-controlled prevalence study. *Spine (Phila Pa 1976)*. 1996;21(15):1737-1744.
56. Bamsley L, Lord SM, Wallis BJ, Bogduk N. The prevalence of chronic cervical zygapophysial joint pain after whiplash. *Spine (Phila Pa 1976)*. 1995;20(1):20-25.
57. Smith AD, Jull G, Schneider G, Frizzell B, Hooper RA, Sterling M. A comparison of physical and psychological features of responders and non-responders to cervical facet blocks in chronic whiplash. *BMC Musculoskelet Disord*. 2013;14:313.
58. Rubinstein SM, Pool JJ, van Tulder MW, Riphagen II, de Vet HC. A systematic review of the diagnostic accuracy of provocative tests of the neck for diagnosing cervical radiculopathy. *Eur Spine J*. 2007;16(3):307-319.
59. Malanga GA, Landes P, Nadler SF. Provocative tests in cervical spine examination: historical basis and scientific analyses. *Pain Physician*. 2003;6(2):199-205.
60. Cook CE, Hegedus E, Pietrobon R, Goode A. A pragmatic neurological screen for patients with suspected cord compressive myelopathy. *Phys Ther*. 2007;87(9):1233-1242.
61. Cohen SP, Bajwa ZH, Kraemer JJ, et al. Factors predicting success and failure for cervical facet radiofrequency denervation: a multi-center analysis. *Reg Anesth Pain Med*. 2007;32(6):495-503.
62. Matsumoto M, Fujimura Y, Suzuki N, et al. MRI of cervical intervertebral discs in asymptomatic subjects. *J Bone Joint Surg Br*. 1998;80(1):19-24.
63. Lehto JJ, Tertti MO, Komu ME, Paajanen HE, Tuominen J, Korhano MJ. Age-related MRI changes at 0.1 T in cervical discs in asymptomatic subjects. *Neuroradiology*. 1994;36(1):49-53.
64. American Association of Electrodiagnostic Medicine. Practice parameter for needle electromyographic evaluation of patients with suspected cervical radiculopathy: summary statement. *Muscle Nerve*. 1999;22(suppl 8):S209-S221.
65. Ashkan K, Johnston P, Moore AJ. A comparison of magnetic resonance imaging and neurophysiological studies in the assessment of cervical radiculopathy. *Br J Neurosurg*. 2002;16(2):146-148.
66. Anderberg L, Annertz M, Rydholm U, Brandt L, Säveland H. Selective diagnostic nerve root block for the evaluation of radicular pain in the multilevel degenerated cervical spine. *Eur Spine J*. 2006;15(6):794-801.
67. Sasso RC, Macadaeg K, Nordmann D, Smith M. Selective nerve root injections can predict surgical outcome for lumbar and cervical radiculopathy: comparison to magnetic resonance imaging. *J Spinal Disord Tech*. 2005;18(6):471-478.

68. Cohen SP, Hurlley RW. The ability of diagnostic spinal injections to predict surgical outcomes. *Anesth Analg*. 2007; 105(6):1756-1775.
69. Kay TM, Gross A, Goldsmith CH, et al. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev*. 2012;8:CD004250.
70. Sihawong R, Janwantanakul P, Sitthipornvorakul E, Pensri P. Exercise therapy for office workers with nonspecific neck pain: a systematic review. *J Manipulative Physiol Ther*. 2011; 34(1):62-71.
71. Bertozzi L, Gardenghi I, Turoni F, et al. Effect of therapeutic exercise on pain and disability in the management of chronic nonspecific neck pain: systematic review and meta-analysis of randomized trials. *Phys Ther*. 2013;93(8): 1026-1036.
72. Kuijper B, Tans JT, Beelen A, Nollet F, de Visser M. Cervical collar or physiotherapy versus wait and see policy for recent onset cervical radiculopathy: randomised trial. *BMJ*. 2009; 339:b3883.
73. Thoomes EJ, Scholten-Peeters W, Koes B, Falla D, Verhagen AP. The effectiveness of conservative treatment for patients with cervical radiculopathy: a systematic review. *Clin J Pain*. 2013;29(12):1073-1086.
74. Furlan AD, Yazdi F, Tsertsvadze A, et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med*. 2012;2012:953139.
75. Hurwitz EL, Carragee EJ, van der Velde G, et al; Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. Treatment of neck pain: noninvasive interventions; results of the Bone and Joint Decade 2000-2010 Task Force on Neck Pain and Its Associated Disorders. *Spine (Phila Pa 1976)*. 2008;33(4, suppl):S123-S152.
76. Kong LJ, Zhan HS, Cheng YW, Yuan WA, Chen B, Fang M. Massage therapy for neck and shoulder pain: a systematic review and meta-analysis. *Evid Based Complement Alternat Med*. 2013;2013:613279.
77. Patel KC, Gross A, Graham N, et al. Massage for mechanical neck disorders. *Cochrane Database Syst Rev*. 2012;9:CD004871.
78. Graham N, Gross A, Goldsmith CH, et al. Mechanical traction for neck pain with or without radiculopathy. *Cochrane Database Syst Rev*. 2008;(3):CD006408.
79. Kroelings P, Gross A, Goldsmith CH, et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev*. 2009;(4):CD004251.
80. Cramer H, Lauche R, Hohmann C, et al. Randomized-controlled trial comparing yoga and home-based exercise for chronic neck pain. *Clin J Pain*. 2013;29(3):216-223.
81. Bronfort G, Evans R, Anderson AV, Svendsen KH, Bracha Y, Grimm RH. Spinal manipulation, medication, or home exercise with advice for acute and subacute neck pain: a randomized trial. *Ann Intern Med*. 2012;156(1, pt 1):1-10.
82. White AP, Arnold PM, Norvell DC, Ecker E, Fehlings MG. Pharmacologic management of chronic low back pain: synthesis of the evidence. *Spine (Phila Pa 1976)*. 2011;36(21, suppl): S131-S143.
83. Flood J. The role of acetaminophen in the treatment of osteoarthritis. *Am J Manag Care*. 2010;16(suppl, Management): S48-S54.
84. Hsieh LF, Hong CZ, Chen SH, Chen CC. Efficacy and side effects of diclofenac patch in treatment of patients with myofascial pain syndrome of the upper trapezius. *J Pain Symptom Manage*. 2010;39(1):116-125.
85. Borenstein DG, Korn S. Efficacy of a low-dose regimen of cyclobenzaprine hydrochloride in acute skeletal muscle spasm: results of two placebo-controlled trials. *Clin Ther*. 2003;25(4): 1056-1073.
86. Berry H, Liyanage SP, Durance RA, Goode JD, Swannell AJ. A double-blind study of benorylate and chlormezanone in musculoskeletal disease. *Rheumatol Rehabil*. 1981;20(1):46-49.
87. Elenbaas JK. Centrally acting oral skeletal muscle relaxants. *Am J Hosp Pharm*. 1980;37(10):1313-1323.
88. Cohen SP, Mullings R, Abdi S. The pharmacologic treatment of muscle pain. *Anesthesiology*. 2004;101(2):495-526.
89. Scott NA, Guo B, Barton PM, Gerwin RD. Trigger point injections for chronic non-malignant musculoskeletal pain: a systematic review. *Pain Med*. 2009;10(1):54-69.
90. Kamanli A, Kaya A, Ardicoglu O, Ozgocmen S, Zengin FO, Bayik Y. Comparison of lidocaine injection, botulinum toxin injection, and dry needling to trigger points in myofascial pain syndrome. *Rheumatol Int*. 2005;25(8):604-611.
91. Soares A, Andriolo RB, Atallah AN, da Silva EM. Botulinum toxin for myofascial pain syndromes in adults. *Cochrane Database Syst Rev*. 2012;4:CD007533.
92. Göbel H, Heinze A, Reichel G, Hefter H, Benecke R; Dysport Myofascial Pain Study Group. Efficacy and safety of a single botulinum type A toxin complex treatment (Dysport) for the relief of upper back myofascial pain syndrome: results from a randomised double-blind placebo-controlled multicentre study. *Pain*. 2006;125(1-2):82-88.
93. Ojala T, Arokoski JP, Partanen J. The effect of small doses of botulinum toxin A on neck-shoulder myofascial pain syndrome: a double-blind, randomized, and controlled crossover trial. *Clin J Pain*. 2006;22(1):90-96.
94. Qerama E, Fuglsang-Frederiksen A, Kasch H, Bach FW, Jensen TS. A double-blind, controlled study of botulinum toxin A in chronic myofascial pain. *Neurology*. 2006;67(2): 241-245.
95. Nicol AL, Wu II, Ferrante FM. Botulinum toxin type A injections for cervical and shoulder girdle myofascial pain using an enriched protocol design. *Anesth Analg*. 2014;118(6): 1326-1335.
96. De Andrés J, Adsua VM, Palmisani S, Villanueva V, López-Alarcón MD. A double-blind, controlled, randomized trial to evaluate the efficacy of botulinum toxin for the treatment of lumbar myofascial pain in humans. *Reg Anesth Pain Med*. 2010;35(3):255-260.
97. Jabbari B. Botulinum toxin A and chronic low back pain: a randomized, double-blind study. *Neurology*. 2001;56(10):1290-1293.
98. Jabbari B, Machado D. Treatment of refractory pain with botulinum toxins—an evidence-based review. *Pain Med*. 2011; 12(11):1594-1606.
99. Anderberg L, Annertz M, Persson L, Brandt L, Säveland H. Transforaminal steroid injections for the treatment of cervical radiculopathy: a prospective and randomised study. *Eur Spine J*. 2007;16(3):321-328.
100. Manchikanti L, Malla Y, Cash KA, McManus CD, Pampati V. Fluoroscopic cervical interlaminar epidural injections in managing chronic pain of cervical postsurgery syndrome: preliminary results of a randomized, double-blind, active control trial. *Pain Physician*. 2012;15(1):13-25.
101. Manchikanti L, Malla Y, Cash KA, McManus CD, Pampati V. Fluoroscopic epidural injections in cervical spinal stenosis: preliminary results of a randomized, double-blind, active control trial. *Pain Physician*. 2012;15(1):E59-E70.
102. Manchikanti L, Cash KA, Pampati V, Wargo BW, Malla Y. A randomized, double-blind, active control trial of fluoroscopic cervical interlaminar epidural injections in chronic pain of cervical disc herniation: results of a 2-year follow-up [published correction appears in *Pain Physician*. 2013;16(6): 609]. *Pain Physician*. 2013;16(5):465-478.
103. Stav A, Ovadia L, Sternberg A, Kaadan M, Weksler N. Cervical epidural steroid injection for cervicobrachialgia. *Acta Anaesthesiol Scand*. 1993;37(6):562-566.
104. Cohen SP, Hayek S, Semenov Y, et al. Epidural steroid injections, conservative treatment or combination treatment for cervical radiculopathy: a multi-center, randomized, comparative-effectiveness study. *Anesthesiology*. 2014; 121(5):1045-1055.

105. Bicket MC, Gupta A, Brown CH IV, Cohen SP. Epidural injections for spinal pain: a systematic review and meta-analysis evaluating the "control" injections in randomized controlled trials. *Anesthesiology*. 2013;119(4):907-931.
106. Manchikanti L, Boswell MV, Singh V, Pampati V, Damron KS, Beyer CD. Prevalence of facet joint pain in chronic spinal pain of cervical, thoracic, and lumbar regions. *BMC Musculoskelet Disord*. 2004;5:15.
107. Lord SM, Bamsley L, Wallis BJ, McDonald GJ, Bogduk N. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. *N Engl J Med*. 1996;335(23):1721-1726.
108. Stovner LJ, Kolstad F, Helde G. Radiofrequency denervation of facet joints C2-C6 in cervicogenic headache: a randomized, double-blind, sham-controlled study. *Cephalalgia*. 2004;24(10):821-830.
109. Kim KH, Choi SH, Kim TK, Shin SW, Kim CH, Kim JI. Cervical facet joint injections in the neck and shoulder pain. *J Korean Med Sci*. 2005;20(4):659-662.
110. Bamsley L, Lord SM, Wallis BJ, Bogduk N. Lack of effect of intraarticular corticosteroids for chronic pain in the cervical zygapophyseal joints. *N Engl J Med*. 1994;330(15):1047-1050.
111. Anderberg L, Annertz M, Brandt L, Säveland H. Selective diagnostic cervical nerve root block—correlation with clinical symptoms and MRI-pathology. *Acta Neurochir (Wien)*. 2004;146(6):559-565.
112. Beynon R, Hawkins J, Laing R, et al. The diagnostic utility and cost-effectiveness of selective nerve root blocks in patients considered for lumbar decompression surgery: a systematic review and economic model. *Health Technol Assess*. 2013;17(19):1-88.
113. Persson LC, Carlsson CA, Carlsson JY. Long-lasting cervical radicular pain managed with surgery, physiotherapy, or a cervical collar: a prospective, randomized study. *Spine (Phila Pa 1976)*. 1997;22(7):751-758.
114. Persson LC, Moritz U, Brandt L, Carlsson CA. Cervical radiculopathy: pain, muscle weakness and sensory loss in patients with cervical radiculopathy treated with surgery, physiotherapy or cervical collar: a prospective, controlled study. *Eur Spine J*. 1997;6(4):256-266.
115. Engquist M, Löfgren H, Öberg B, et al. Surgery versus nonsurgical treatment of cervical radiculopathy: a prospective, randomized study comparing surgery plus physiotherapy with physiotherapy alone with a 2-year follow-up. *Spine (Phila Pa 1976)*. 2013;38(20):1715-1722.
116. Cesaroni A, Nardi PV. Plasma disc decompression for contained cervical disc herniation: a randomized, controlled trial. *Eur Spine J*. 2010;19(3):477-486.
117. Kadanka Z, Bednarik J, Vohánka S, et al. Conservative treatment versus surgery in spondylotic cervical myelopathy: a prospective randomised study. *Eur Spine J*. 2000;9(6):538-544.
118. Kadaňka Z, Bednařík J, Novotný O, Urbánek I, Dušek L. Cervical spondylotic myelopathy: conservative versus surgical treatment after 10 years. *Eur Spine J*. 2011;20(9):1533-1538.
119. Kadanka Z, Mares M, Bednarik J, et al. Predictive factors for spondylotic cervical myelopathy treated conservatively or surgically. *Eur J Neurol*. 2005;12(1):55-63.
120. Jacobs W, Van der Gaag NA, Tuschel A, et al. Total disc replacement for chronic back pain in the presence of disc degeneration. *Cochrane Database Syst Rev*. 2012;9:CD008326.
121. Brox JI, Nygaard ØP, Holm I, Keller A, Ingebrigtsen T, Reikerås O. Four-year follow-up of surgical versus non-surgical therapy for chronic low back pain. *Ann Rheum Dis*. 2010;69(9):1643-1648.
122. Fallah A, Akl EA, Ebrahim S, et al. Anterior cervical discectomy with arthroplasty versus arthrodesis for single-level cervical spondylosis: a systematic review and meta-analysis. *PLoS One*. 2012;7(8):e43407.
123. Yu L, Song Y, Yang X, Lv C. Systematic review and meta-analysis of randomized controlled trials: comparison of total disc replacement with anterior cervical decompression and fusion. *Orthopedics*. 2011;34(10):e651-e658.
124. Peolsson A, Peolsson M. Predictive factors for long-term outcome of anterior cervical decompression and fusion: a multivariate data analysis. *Eur Spine J*. 2008;17(3):406-414.