


People with “rage” disorder twice as likely to have a latent toxoplasmosis parasite infection

Posted on March 23, 2016 by [Kevin Jiang](#) in [At the Bench](#), [Don't Miss](#)



(<https://sciencelife.uchospitals.edu/2014/07/22/what-does-it-mean-when-2-billion-people-share-their-brain-with-a-parasite/8-blue/>)

The protozoan *Toxoplasma gondii*, tissue cyst in brain (Photo: D. Ferguson, Oxford University)

Individuals with a psychiatric disorder involving recurrent bouts of extreme, impulsive anger—road rage, for example—are more than twice as likely to have been exposed to a common parasite than healthy individuals with no psychiatric diagnosis.

In a study involving 358 adult subjects, a team led by researchers from the University of Chicago found that toxoplasmosis, a relatively harmless parasitic infection carried by an estimated 30 percent of all humans, is associated with intermittent explosive disorder and increased aggression.

The findings are published in the *Journal of Clinical Psychiatry* (<http://dx.doi.org/10.4088/JCP.14m09621>) on March 23, 2016.


“Our work suggests that latent infection with the *toxoplasma gondii* parasite may change brain chemistry in a fashion that increases the risk of aggressive behavior,” said senior study author [Emil Coccaro](http://cnpru.bsd.uchicago.edu/coccaro.html) (<http://cnpru.bsd.uchicago.edu/coccaro.html>), MD, Ellen. C. Manning Professor and Chair of Psychiatry

and Behavioral Neuroscience at the University of Chicago.

"However, we do not know if this relationship is causal, and not everyone that tests positive for toxoplasmosis will have aggression issues," Coccaro said, adding that additional studies are needed.

Intermittent explosive disorder (<http://www.mayoclinic.org/diseases-conditions/intermittent-explosive-disorder/basics/definition/con-20024309>) (IED) is defined by the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition, as recurrent, impulsive, problematic outbursts of verbal or physical aggression that are disproportionate to the situations that trigger them. IED is thought to affect as many as 16 million Americans, more than bipolar disorder and schizophrenia combined.

As part of their pioneering research to improve diagnosis and treatment for IED and impulsive aggression, Coccaro and his colleagues examined possible connections to toxoplasmosis (<http://www.cdc.gov/parasites/toxoplasmosis/>), an extremely common parasitic infection. Transmitted through the feces of infected cats, undercooked meat or contaminated water, toxoplasmosis is typically latent and harmless for healthy adults. However, it is known to reside in brain tissue, and has been linked to several psychiatric diseases, including schizophrenia, bipolar disorder and suicidal behavior.



(<https://sciencelife.uchospitals.edu/2016/03/23/people-with-rage-are-disorder-twice-as-likely-to-have-a-latent-toxoplasmosis-parasite-infection/toxoplasma-gondii-life-cycle/>)

Toxoplasmosis life cycle. Courtesy CDC.

The research team recruited 358 adult subjects from the U.S., who were evaluated for IED, personality disorder, depression and other psychiatric disorders. Study participants were also scored on traits including anger, aggression and impulsivity. Participants fell into one of three groups. Roughly one third had IED. One third were healthy controls with no psychiatric history. The remaining third were individuals diagnosed with some psychiatric disorder, but not IED. This last group served as a control to distinguish IED from possible confounding psychiatric factors.

Hold your cats

The research team found that IED-diagnosed group was more than twice as likely to test positive for toxoplasmosis exposure (22 percent) as measured by a blood test, compared to the healthy control group (9 percent).

Around 16 percent of the psychiatric control group tested positive for toxoplasmosis, but had similar aggression and impulsivity scores to the healthy control group. IED-diagnosed subjects scored much higher on both measures than either control group.

Across all study subjects, toxoplasmosis-positive individuals scored significantly higher on scores of anger and aggression. The team noted a link between toxoplasmosis and increased impulsivity, but when adjusted for aggression scores, this link became non-significant. This finding suggests toxoplasmosis and aggression are most strongly correlated.

However, the authors caution that the study results do not address whether toxoplasmosis infection may cause increased aggression or IED.

“Correlation is not causation, and this is definitely not a sign that people should get rid of their cats,” said study co-author [Royce Lee \(https://psychiatry.uchicago.edu/directory/royce-lee-md\)](https://psychiatry.uchicago.edu/directory/royce-lee-md), MD, Associate Professor of Psychiatry and Behavioral Neuroscience at the University of Chicago. “We don’t understand the mechanisms involved—it could be an increased inflammatory response, direct brain modulation by the parasite, or even reverse causation where aggressive individuals tend to have more cats or eat more undercooked meat. Our study signals the need for more research and more evidence in humans.”

Coccaro and his team are now further examining the relationship between toxoplasmosis, aggression and IED. If better understood, this connection may inform new strategies to diagnose or treat IED in the future.

“It will take experimental studies to see if treating a latent toxoplasmosis infection with medication reduces aggressiveness,” Coccaro said. “If we can learn more, it could provide rationale to treat IED in toxoplasmosis-positive patients by first treating the latent infection.”

##

The study, “*Toxoplasma gondii* Infection: Relationship With Aggression in Psychiatric Subjects,” was supported by the National Institute of Mental Health (RO1 MH60836, RO1 MH63262, RO1 MH66984), the American Foundation for Suicide Prevention, and the Veterans Integrated Service Network 19 Mental Illness Research, Education and Clinical Center. Additional authors include Maureen W. Groer, Adem Can, Mary Coussons-Read and Teodor T. Postolache.

- Agresion
- Anger
- behavioral neuroscience
- Emil Coccaro
- Imulsitivty
- intermittent explosive disorder
- Parasite infection
- Psychiatry
- Royce Lee
- toxoplasmosis

About Kevin Jiang ([147 Articles](#))

Kevin Jiang is a Science Writer and Media Relations Specialist at the University of Chicago Medicine. He focuses on neuroscience and neurosurgery, orthopedics, psychology, genetics, biology, evolution, biomedical and basic science research.

Connect With Us

Follow Science Life via Email

For Patients

UChicago Links

Copyright © 2016 | [Blog at WordPress.com.](#) | [The MH Magazine Theme.](#)